

AD-A235 577



PL-TR-91-2028

2

Seismic Event Location at Regional Distances

C. Thurber

Research Foundation of State University  
of New York  
P. O. Box 9  
Albany, NY 12201

22 March 1991

Final Report  
22 July 1988 - 14 January 1991

Approved for public release; distribution unlimited

DTIC  
ELECTE  
MAY 14 1991  
S B D



PHILLIPS LABORATORY  
AIR FORCE SYSTEMS COMMAND  
HANSCOM AIR FORCE BASE, MASSACHUSETTS 01731-5000

NOT A COPY


91 5 14 026


SPONSORED BY  
Defense Advanced Research Projects Agency  
Nuclear Monitoring Research Office  
ARPA ORDER NO. 5299

MONITORED BY  
Phillips Laboratory  
F19628-88-K-0037

The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U.S. Government.

This technical report has been reviewed and is approved for publication.

  
JAMES F. LEWKOWICZ  
Contract Manager  
Solid Earth Geophysics Branch  
Earth Sciences Division

  
JAMES F. LEWKOWICZ  
Branch Chief  
Solid Earth Geophysics Branch  
Earth Sciences Division

FOR THE COMMANDER

  
FOR: DONALD H. ECKHARDT, Director  
Earth Sciences Division

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.


If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify PL/IMA, Hanscom AFB, MA 01731-5000. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 22 March 1991		3. REPORT TYPE AND DATES COVERED Final 22 Jul 88-14 Jan 91
4. TITLE AND SUBTITLE  Seismic Event Location at Regional Distances			5. FUNDING NUMBERS  F19628-88-K-0037 PR 8A10 TA DA WU AI	
6. AUTHOR(S)  C. Thurber				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Research Foundation of SUNY P.O. Box 9 Albany, NY 12201			8. PERFORMING ORGANIZATION REPORT NUMBER  431-6722A-2	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Phillips Laboratory Hanscom AFB, MA 01731-5000  DARPA 1400 Wilson Blvd Arlington, VA 22209-2308  Contract Manager: James Lewkowicz/LWH			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  PL-TR-91-2028	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; Distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  Three-component data from regional seismic events recorded by the former NRDC-Soviet Academy of Sciences regional seismic network in Kazakhstan, USSR, have been analyzed with the primary goal of improving regional seismic event location capability. Data from these events were used in the investigation of the following problems related to regional event location: 1) determination of wave arrival azimuth; 2) observability and value of secondary phase arrivals; 3) evaluation and improvement of regional event location algorithms; 4) independent determination of "master event" locations. Starting with very little prior information, we have demonstrated the potential for a sparse seismic network of three 3-component stations to locate events over a wide region with reasonable accuracy and precision, both for epicenter and depth. Our basic findings are: that arrival azimuth can be determined with reasonable precision, but the data provide little in the way of location constraint in most cases; secondary phase arrivals are routinely observable over a wide distance range, and they provide important location constraints; existing location algorithms perform well and provide appropriate estimates of location uncertainty, but must be modified for far-regional applications; numerous "master events" have been identified and utilized to improve our location capability.				
14. SUBJECT TERMS  Event location, regional seismic waves, satellite imagery, depth determination			15. NUMBER OF PAGES 40	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT  UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE  UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT  UNCLASSIFIED	20. LIMITATION OF ABSTRACT  SAR	

## TABLE OF CONTENTS

	<u>PAGE</u>
1. Summary	1
2. Accomplishments	2
2.1. Task Objectives	2
2.2. Technical Problem	2
2.3. General Methodology	3
2.4. Technical Results	9
3. Conclusions and Recommendations	21
4. References	22



<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution/ _____	
<b>Availability Codes</b>	
Dist	Avail and/or Special
A-1	

## 1. Summary

We have carried out a detailed analysis of three-component data from regional seismic events recorded by the former NRDC-Soviet Academy of Sciences regional seismic network in Kazakhstan, USSR, with the primary goal of improving regional seismic event location capability. The data from these events were used in the investigation of the following problems related to regional event location: 1) determination of wave arrival azimuth; 2) observability and value of secondary phase arrivals; 3) evaluation and improvement of regional event location algorithms; 4) independent determination of "master event" locations. Starting with very little prior information, we have demonstrated the potential for a sparse seismic network of three 3-component stations, with an aperture of about 400 km, to locate events over a wide region with reasonable accuracy and precision, both for epicenter and depth. Our basic findings are: that arrival azimuth can be determined with reasonable precision, but the data provide little in the way of location constraint in most cases; secondary phase arrivals are routinely observable over a wide distance range, and they provide important location constraints; existing location algorithms perform well and provide appropriate estimates of location uncertainty, but must be modified for far-regional applications; numerous "master events" have been identified and utilized to improve our location capability. Some keys to the success of this effort have been: (1) the initial availability of an adequate crustal model, coupled with the inferred modest level of lateral variations in crustal structure within the array; (2) the availability of data from known sources (the 1987 chemical explosions) to aid the calibration of travel time models; (3) the existence of identifiable secondary P arrivals; (4) the availability of satellite imagery to confirm the source location of some events; (5) the ability to pursue a multifaceted approach to the investigation of the overall location problem (empirical, theoretical, observational). We can make several specific recommendations for future efforts related to regional event location: (1) employ array studies to help identify and characterize secondary regional phases; (2) use short time windows at the onset of the P arrival for precise azimuth estimation; (3) obtain total coverage of the region in question by satellite imagery - including multiple sources of data (e.g., SPOT, LANDSAT). Overall, we regard regional seismic event location to be a difficult but quite tractable problem.

## 2. Accomplishments

### 2.1. Task Objectives

We have carried out a detailed analysis of three-component data from regional seismic events recorded by the former NRDC-Soviet Academy of Sciences regional seismic network in Kazakhstan, USSR, with the primary goal of improving regional seismic event location capability. One basic task was the cataloging of locations for some of the regional events detected by the network. The data from these events were used in the investigation of the following problems related to regional event location: 1) determination of wave arrival azimuth; 2) observability and value of secondary phase arrivals; 3) evaluation and improvement of regional event location algorithms; 4) independent determination of "master event" locations. An improved model for the regional seismic velocity structure was sought in conjunction with the above tasks.

### 2.2. Technical Problem

Seismic event location remains a fundamental component of monitoring efforts related to verifying nuclear test ban or test limitation treaties. Event location is important both as a basis for discrimination by itself and as a starting point for the analysis of wave propagation and attenuation. The earthquake location problem is relatively well understood on a theoretical basis (Thurber, 1986). However, it can be expected that event location will be a non-trivial problem for in-country regional networks like those considered in the recent literature (e.g., Evernden et al., 1986).

Valuable experience in the study of regional wave propagation in the U.S. has been gained from the operation of the Regional Seismic Test Network (RSTN), but the direct applicability of that experience to the regional monitoring of weapons testing in the U.S.S.R. is questionable, due to significant differences in crustal structure and attenuation characteristics. However it is now possible to obtain high-quality digital data from stations within the Soviet Union. An agreement between the National Resources Defense Council, Inc., and the Academy of Sciences of the U.S.S.R. led to the establishment of a three-station seismic network in the spring of 1987 in eastern Kazakhstan in the Soviet Union. Each station consisted of several sets of 3-component instruments, recorded digitally at 250 samples per second per channel using a triggered system (Berger et al., 1988). As part of the agreement, a similar network was set up in the western U.S. The stations of the two networks encircled the Kazakhstan and Nevada nuclear test sites (KTS and NTS), respectively, at distances of about 200 km. The stated purpose of these networks was to collect data relevant to seismic monitoring of nuclear weapons tests (Berger et al., 1987). In late

1988, the Kazakh network was dismantled, and new stations were set up (at much greater distances from KTS) through an agreement established by with the Incorporated Research Institutions for Seismology (IRIS) and the U. S. Geological Survey.

Due to the sparseness of the Kazakhstan network, most regional events (predominantly mining blasts) did not trigger all three stations, and often triggered only one. Similar detection difficulties can be expected for any sparse regional monitoring network. Standard earthquake location algorithms that use P and S wave arrival times alone cannot be expected to yield satisfactory results under such conditions, particularly for small to moderate sized events. Alternative methods and/or additional data are needed for adequate constraint of event locations.

### 2.3. General Methodology

A promising approach to the problem of locating seismic events with a sparse regional network is the use of arrival times from multiple regional phases ( $P_g$ ,  $L_g$  and  $R_g$ , and possibly  $P^*$ ,  $S^*$ , in addition to  $P_n$ ,  $S_n$ ) and arrival azimuths. Bratt and Bache (1988) describe an earthquake location algorithm that uses arrival times and azimuths to estimate regional event locations. Secondary arrivals are handled computationally exactly the same as first arrivals. Arrival azimuth information can be incorporated by adding azimuth residuals to the residual vector and the corresponding rows of partial derivatives to the Jacobean (Bratt and Bache, 1988). The partial derivatives for azimuth are derived directly from the source-receiver geometry. Bratt and Bache (1988) applied their algorithm successfully to array data from NORESS and FINESA for events at distance ranges of 200 to 1500 km. Magotra et al. (1987) and Ruud et al. (1988) describe single-station approaches using arrival times and azimuths (slowness vectors) that are conceptually similar to the Bratt and Bache approach. They applied their methods to data from single-site RSTN and NORESS three-component records, respectively.

The standard earthquake location method iteratively solves a matrix equation relating hypocenter adjustments to arrival time residuals via the Jacobian matrix, consisting of the partial derivatives of arrival time with respect to the event coordinates and origin time (Thurber, 1986), with the iterations stopping when some convergence criterion is reached. Arrival azimuth information can be incorporated as additional information for determining the location by adding azimuth residuals to the residual vector and the corresponding rows of partial derivatives to the Jacobian matrix (Bratt and Bache, 1988). The partial derivatives for azimuth are derived directly from the source-receiver geometry. Both the event depth and origin time are independent of the azimuth. In the algorithm TTAZLOC (Bratt and Bache, 1988), the final solution is obtained using iterative damped least squares. The estimate of location uncertainty is derived using a combination


of *a priori* and *a posteriori* data uncertainties (Jordan and Sverdrup, 1981). The *a priori* uncertainty can incorporate estimated uncertainties due to measurement error and inexact knowledge of velocity structure (Bratt and Bache, 1988; Pavlis, 1986). In this initial investigation, we use the estimated uncertainties in arrival time reading and arrival azimuth determination for the *a priori* values. The former typically are 0.25 s for P and 0.5 s for S arrivals, based on subjective estimation of reading quality, while the latter are on the order of 5°, based on the standard deviation of the computed arrival azimuth within the selected window. We also assume a value for the Bayesian parameter K of 8 (Jordan and Sverdrup, 1981). A value for K of 8 assigns 8 degrees of freedom to the *a priori* uncertainty, and implies an expected standard deviation of the *a posteriori* standard deviation variable (the reciprocal of the normalized *a posteriori* uncertainty) of 25% about a mean of 1.0 (Jordan and Sverdrup, 1981). Note that in the discussion and tables, all error ellipses will be represented by the major axes of the 90% confidence ellipses.

A crustal model is required to compute travel times of the various phases. Fortunately, Soviet Deep-Seismic-Sounding (DSS) surveys have been carried out in the region, including one line just to the west of Karkaralinsk (Figure 1), yielding estimates of crust and upper mantle P velocities and crustal thickness. Results from these profiles are summarized by Belyaevsky et al. (1973), Antonenko (1984), and Leith (1987). Crustal thickness varies between about 45 and 55 kilometers in the immediate vicinity of the network (Belyaevsky et al., 1973). We have adopted a layered approximation to the P-wave velocity model reported by Antonenko (1984), shown in Figure 2, modified to account for the low-velocity granites that underly the station sites (Leith, 1987). The model predicts a  $P_n$  crossover distance of about 220 km.


There is little published information on the S velocity structure in the region. Priestley et al. (1988) carried out a preliminary teleseismic waveform inversion for records at stations BAY and KKL using the method of Owens et al. (1984), deriving a layered model for the S velocity structure beneath those stations. For comparison, taking the DSS profile results and assuming a constant  $V_p/V_s$  ratio of 1.73 gives values for  $V_s$  that are consistent with the teleseismic waveform analysis of Priestley et al. (1988) for the upper crust, but somewhat lower than Priestley's for the lower crust. This is in agreement with the finding of Alekseev et al. (1988) that  $V_p/V_s$  is higher in the upper crust than the lower crust in the area of Kazakhstan near the Tien Shan. Given the uncertainties, however, we have chosen to calculate the event locations using two different S velocity models (Table 1), one with constant  $V_p/V_s$  ratio (Model A), and another using the model of Priestley et al. (1988) (Model B).

The location algorithm TTAZLOC is structured to be able to utilize arrival times of secondary phases, assuming they can be identified and modeled correctly. Travel time calculations





**Figure 1.** Configuration of the NRDC-Soviet Academy seismic network, located near the Kazakhstan Test Site (KTS) on the steppes of Central Asia. Stations BAY, KKL, and KSU are indicated by the solid triangles. Also shown are the sites of the September 1987 chemical explosions (+) and the location of a Deep-Seismic-Sounding profile (DSS).



**Figure 2.** Comparison of the 1-D velocity model for the Kazakhstan area from Deep Seismic Sounding (Antonenko, 1984), and its modifications based on empirical location work, mostly based on the 1987 chemical explosion (Thurber et al., 1989) and synthetic modeling of regional seismograms (this work).

Table 1

Velocity models used for event location

MODEL A

Depth range (km)	Vp (km/sec)	Vs (km/sec)
0 - 5	5.40	3.05
5 - 10	6.15	3.50
10 - 20	6.35	3.60
20 - 30	6.55	3.70
30 - 40	6.75	3.85
40 - 50	6.95	3.95
50 +	8.20	4.65

MODEL B

Depth range (km)	Vp (km/sec)	Vs (km/sec)
0 - 5	5.40	3.30
5 - 10	6.15	3.40
10 - 20	6.35	3.50
20 - 30	6.55	3.70
30 - 40	6.75	4.10
40 - 50	6.95	4.30
50 +	8.20	4.70

for the P velocity model in Figure 2 indicate that an upper-crustal refraction along the 6.35 km/sec layer at 10 km depth gives the first arrival from surface sources in the distance range of about 100 to 220 km, and also suggests it might produce a significant secondary arrival beyond that distance. We will denote this phase as  $P_g$ , following the notation of Aki and Richards (1980; p. 213). The phases we use for locations include  $P_n$ ,  $P_g$ ,  $S_n$ , and  $S_g$ , when observable; we have not used  $L_g$ , as we regard it as having a less precisely measurable arrival time.

For seismic arrays or sparse networks, the azimuth of arriving phases can provide crucial constraints for determining event locations. We determine arrival azimuths using a method similar to that described by Magotra et al. (1987) for estimating the polarization direction of arriving seismic phases. For a polarized signal in the presence of noise, the eigenvector corresponding to the largest eigenvalue of the covariance matrix for the signal components gives the direction of polarization, and the ratio of eigenvalues measures the rectilinearity of particle motion (Kanasewich, 1981). Since we are interested just in the arrival azimuth, the horizontal component seismograms from a station are windowed (usually over 100 to 250 samples or 0.4 to 1.0 s) and demeaned, and the 2-by-2 signal covariance matrix  $C$  is computed:

$$C = \begin{bmatrix} \text{Var}[NS] & \text{Cov}[NS,EW] \\ \text{Cov}[NS,EW] & \text{Var}[EW] \end{bmatrix}$$

where NS and EW represent the north-south and east-west component time series, respectively. From the eigenvector for the largest eigenvalue of  $C$ ,  $E_{\lambda_{\max}} = [e_1 \ e_2]$  the polarization direction  $\phi$  for the time window can be computed from  $\tan \phi = e_2 / e_1$ . For the P-wave,  $\phi$  will be the apparent back-azimuth to the event. The inherent  $180^\circ$  ambiguity in azimuth can be resolved by using all three components of particle motion (Magotra et al., 1987). We find we are consistently able to determine azimuth estimates from the first P arrival, usually with an estimated uncertainty of  $5^\circ$  or less. Particle motion plots are also examined for a simple check of the results.

Our methodology for evaluating the utility of secondary arrivals and arrival azimuths for the purpose of regional seismic event location has been to approach the issue from several complementary directions to seek self-consistent results. These directions are observational, empirical, and theoretical. Observationally, we have examined and analyzed regional seismic data from the Kazakhstan area to investigate the quality of secondary phase and arrival azimuth data. Empirically, we have assessed the ability to determine accurate event locations in cases where the true event location is known independently or can be inferred, for example from satellite images. Theoretically, we have evaluated extensively the formal regional event location uncertainty using secondary phase and arrival azimuth data in conjunction with the traditional first arrival data. Overall, we feel we have developed a consistent picture of regional event location capability, but

one which leaves room for improvement as our data and our understanding of regional wave propagation improves.

## 2.4. Technical Results

The project duration was approximately 30 months, 7/22/88 to 12/31/90. Our efforts during the period were divided among four components of study: arrival azimuth, secondary arrivals, location algorithm, and master events. Progress achieved in these four project components is described in the following sections.

### 2.4.1. Arrival Azimuth

A number of investigators have examined methods for arrival azimuth determination from 3-component seismic data (Magotra et al., 1987; Jurkevics, 1988; Christoffersson et al., 1988). We have found the principal components analysis method to be quite successful, particularly for broadband data analyzed in short time windows. Figure 3 shows an example of arrival azimuth estimation using principal components analysis. The data are from the 1987 chemical explosion 1 in Kazakhstan recorded at station KKL (Given et al., 1990). The arrival azimuth and rectilinearity are calculated from the horizontal components, using a sliding time window of 0.4 seconds. No filtering (Jurkevics, 1988) or time-lagged weighting (Magotra et al., 1987) is applied. Note the significant increase in rectilinearity and stability of the azimuth estimate associated with the regional seismic phases identified in the seismogram. A reanalysis of several other selected events from the NRDC data set provides clear justification for the use of short, early windows in the P waveform for computing arrival azimuth. This strongly suggests that automated algorithms for estimating arrival azimuth should use one or more polarization strength measures to select an appropriate window for computing an azimuth estimate.

A comparison was also made between time-domain (principal components) and frequency-domain (multiple spectral taper) polarization techniques for determining arrival azimuth from the chemical explosion data. Our work has shown that the two methods obtain compatible results for matching time windows. However, the time-domain method has the advantage of finer time resolution, allowing identification of scattered and/or converted waves in the coda of the desired direct arrival. The frequency-domain method is useful for indicating the bandwidth over which reliable azimuth information is present (usually 1 to 15 Hz).

Two critical questions are (1) what are the accuracy and precision of arrival azimuth estimates, and (2) how useful is arrival azimuth in constraining regional event locations. We have




Figure 3. Arrival azimuth estimation using principal components analysis. The top panel shows 5 seconds of the 3-component seismograms from Chemical Explosion 1 recorded at station KKL, while the bottom panel shows the calculated arrival azimuth and rectilinearity of the horizontal components, using a sliding time window of 0.4 seconds.

Table 2

Estimated and true arrival azimuths for 1987 Kazakhstan chemical explosions 1 and 2

Explosion		BAY	KKL	KSU
1	Estimated ( $\pm 1 \sigma$ )	$255 \pm 2$	$293 \pm 1$	$276 \pm 2$
	True	257	296	277
	Error	2	3	1
2	Estimated ( $\pm 1 \sigma$ )	$121 \pm 2$	$56 \pm 2$	$279 \pm 4$
	True	125	61	272
	Error	4	5	7

---

limited answers to the first question, and a fairly complete answer to the second. Our analysis of data from the 1987 chemical explosions in Kazakhstan suggests that azimuth accuracy is on the order of  $5^\circ$  and that precision is no better than  $2^\circ$ , on average (see Table 2 above). However, the latter estimate is not based on a formal error assessment, rather on an evaluation of azimuth stability within the selected time window. Other studies have suggested that precision is only 5 to  $10^\circ$ .

Regarding constraining regional event locations, our empirical and theoretical analyses demonstrate the limited utility of azimuth data. For single-station locations, of course, arrival azimuth provides the only information about source direction, and thus is critical. However, only very poor constraint on epicenter is obtainable if the azimuth uncertainty is on the order of  $5^\circ$ ; uncertainty increases roughly linearly with epicentral distance, at a rate of about 0.1 km per km. Thus single-station location uncertainty will be quite high at regional distances (200 to 2000 km, say). This situation can be improved significantly only if the uncertainty in azimuth estimation can be reduced substantially, for example with the use of arrays and the correction for systematic errors.

For two-station locations, azimuth data contribute only to a slight reduction in epicenter uncertainty, specifically along the line connecting the two observing stations, and contribute nothing to reducing depth uncertainty. We do note, however, that without azimuth data, there is a fundamental ambiguity in source location in this case - two points symmetrically located about the line connecting the two observing stations will always fit the arrival time data equally well. Azimuth information can readily remove this ambiguity. Thus they are essential for providing

directional information but of themselves do not improve two-station location constraint significantly. We have also determined that arrival azimuth data contribute no significant information in the case of three-station locations.

#### 2.4.2. Secondary Arrivals

One of our fundamental hypotheses at the outset of this study was that secondary arrivals could prove to be extremely useful for constraining regional event locations. We have undertaken a three-pronged approach to the evaluation of this hypothesis: empirical (location results using secondary arrivals), theoretical (quantification of hypothetical location capability with secondary arrivals), and observational (observability of secondary arrivals). Our results appear to be self-consistent and very encouraging regarding the utility of secondary arrivals.

Our most significant empirical results concern the determination of source depth with travel time data. A thorough analysis of data from the 1987 chemical explosions in Kazakhstan has shown that source depth can be adequately constrained even with data from a single station if multiple arrivals are used. We relocated chemical explosions 1 and 2 first with depth fixed at 5 km and then with depth free, using velocity model A. For both events, the fixed 5 km depth solutions were notably worse than the original fixed 0 km depth solutions: the mean absolute arrival time residuals increased from 0.08 sec to 0.87 sec for explosion 1 and from 0.24 sec to 0.68 sec for explosion 2. Furthermore, for the solutions with focal depth left free, the final calculated location was in fact at 0 km depth in each case, with estimated focal depth uncertainties of 0.6 km and 1.3 km for explosions 1 and 2, respectively. If we further eliminate the data from station KSU, the location quality remains essentially unaltered for explosion 1, but degrades significantly for explosion 2. In the latter case, constraint on source depth is completely lost. With data from only a single station (either BAY or KKL), the solution for explosion 1 is still stable and reasonably accurate, falling within 8 km in epicenter and 1 km in depth. We attribute these surprisingly successful results for explosion 1 to the availability of multiple secondary arrivals ( $P_g$  and  $S_n$ ). These stations lie just beyond the crossover distance, where  $P_n$  is the first arrival and  $P_g$  can be clearly observed following  $P_n$  (see Figure 3 for an example). Thus we would agree with the claim of Ruud et al. (1988) that focal depth can be determined from data at a single station. A far more thorough analysis will be required to establish the conditions under which depths of regional events can be adequately constrained in general.

By making use of inverse theory, it is possible to analyze the expected location uncertainty and stability (including depth) and evaluate the importance of each arrival time datum as functions of actual event location, given the station locations and observed phases. Our theoretical analysis of location capability (Li and Thurber, 1991) applied Singular Value Analysis (Lawson and



Hanson, 1974) to suites of hypothetical arrival time and azimuth observations of P and S phases at the three NRDC network stations (or subsets thereof). Hypothetical sources were distributed over a region 1000 km by 1000 km, with a grid size of 20 km. Source depths were varied between 0 km and 8 km; we find that location capability is quite insensitive to source depth over this range. We calculated the theoretical location uncertainty (assuming reasonable data variance values), and also the data importance, to assess the solution quality for the various suites of data (stations and phases). Arrival time uncertainty of 0.5 s and arrival azimuth uncertainty of 5° was assumed.

Our theoretical results indicate that source depth can be constrained quite well with adequate regional travel time data, with epicenter somewhat less well constrained. First P and first S arrivals alone at the three stations are not adequate for acceptable epicenter and depth constraint. Location uncertainties for both parameters exceed 30 km for half of the study region. However, adding a single secondary P observation, such as PmP, at all three stations results in depth uncertainties less than 10 km and epicenter uncertainties less than 15 km over more than 95% of the region. Figures 4 and 5 provide a summary of theoretical location uncertainty (depth and epicenter) for various combinations of regional phases.


We also carried out the task of evaluating secondary P phase observability from previously studied events recorded by the former NRDC network surrounding the Soviet KTS. PmP is the most readily observed secondary phase over a substantial distance range, perhaps 125 to 300 km. Pg as a secondary arrival probably has only a very limited range of use just beyond the Pn crossover distance, around 230 to 270 km. We can adequately model the arrival times of these phases (Table 3) using only a slight modification of the model adopted in our previous work on event location (Figure 2 above).

---

Table 3

Misfit of observed P phase times to one-dimensional velocity model

Phase	Mean (sec)	Standard Deviation (sec)
Pg	-0.02	0.4
Pn	+0.05	0.5
PmP	+0.24	0.3



**Figure 4.** Range (box) and mean (vertical bar) of calculated location uncertainty for event depth for suites of possible regional phase observations at 3 stations.




Figure 5. Range (box) and mean (vertical bar) of calculated location uncertainty for event epicenter for suites of possible regional phase observations at 3 stations.

### 2.4.3. Location Algorithm

A comparison was made between travel time calculations in a flat-layered versus spherical-shell earth model to define the limitations for use of a flat-layered model in regional travel time calculations. Our calculations indicate that the travel time discrepancy increases approximately linearly with distance, increasing from 0.5 seconds at 600 km to 1.8 seconds at 2000 km. Thus it will be essential to determine far-regional event locations using a spherical earth model.

Our previous work on regional event location (Thurber et al., 1989) utilized the location algorithm TTAZLOC, which allows the use of "a priori" information in the assessment of event location uncertainty. For our initial work, it was assumed that errors in determining body wave arrival times were the sole contributor to location uncertainty (data-based uncertainty). We are now examining the self-consistency of our a priori uncertainty estimates with the a posteriori uncertainty estimates. The Bayesian statistics approach used in TTAZLOC apportions the contributions to the a posteriori uncertainty estimates according to a "K-weight": a priori uncertainty is treated as providing K data, and residuals from arrival time and azimuth data provide an additional N observational data. Previous studies have used a K-weight of 8, which would be statistically consistent with a normalized standard deviation of the distribution of the standard deviation variable (the reciprocal of the standard deviation) of 0.25, and of course a normalized mean of 1.0 (Jordan and Sverdrup, 1981). For our previous work assuming the data-based uncertainty values (case a), the standard deviation is 0.24 but the mean is 0.81, 20% too low. If we instead assume fixed uncertainty values (in case b of 0.25 seconds for P, 0.5 seconds for S, in case c of 0.5 seconds for P, 1.0 seconds for S, and 5° for azimuth in both cases) reflecting the assumption that inaccuracies in the crustal velocity structure, rather than the data, control location uncertainty, then we produce distributions with standard deviations of 0.19 and 0.17 and means of 0.97 and 1.04 (cases b and c), very close to the correct mean of 1.0 but now below the self-consistent standard deviation value of 0.25. The main implication is that the study of NRDC data by Thurber et al. (1989) underestimated the location uncertainties of regional events by about a factor of 1.5 to 2, given the present state of knowledge of crustal structure. However, a modest improvement in our ability to predict regional seismic wave travel times (P to 0.25 sec, S to 0.5 sec, for example), would reduce the location uncertainties back to quite low levels.

### 2.4.4. Master Events

One of the keys to improving event location capability is the availability of "master events" that can be used for "calibration" of travel time models (or equivalently seismic velocity structure

models). We have used a combination of events with independently known location (chemical and nuclear explosions) and events whose source can be identified via satellite images to produce a set of such master events for the NRDC network. Unfortunately, both types of events are quite limited in number, due to the small number of supervised explosions in Kazakhstan to date (four, two at the same site) and the sparsity of high-quality satellite observations available. The chemical explosions proved particularly valuable for a first-order evaluation of the velocity model used for event location (Thurber et al., 1989).

On September 2 and 3, 1987, three large (10 to 20 ton) chemical explosions were detonated in the vicinity of the network, two at the same location to the northwest of Karaganda and one on the western boundary of the Kazakh test site (Eissler et al., 1987). Their locations are indicated in Figure 1. Eissler et al. (1987) described the characteristics of the seismograms and presented basic analyses of their spectra and size. All three blasts were successfully recorded by each of the network stations. However, we have concentrated our analysis on the data for the first two explosions, as records of the third overlapped a teleseismic arrival from a magnitude 7.2 earthquake in the Macquarie Islands. The event depths were fixed at the surface for the initial set of calculations. In the following discussion, it should be kept in mind that the "true" explosion locations may in fact be somewhat in error, perhaps by as much as a few kilometers, as it is not certain how accurate the maps are that were used by the Soviets to provide the locations.

Explosion 1 was located well outside the network, at sufficient epicentral distance from all the stations for  $P_n$  and  $S_n$  to be clear first arrivals. A strong crustal refracted phase ( $P_g$ ) is also observed at stations BAY and KKL, both about 250 km from the shot site. Despite the unfavorable location with respect to the network, the availability of data at three stations, including numerous secondary arrivals, removes the necessity of incorporating azimuthal information to yield a stable location.

The location estimates for explosion 1 using the two S velocity models are listed in Table 4. The constant  $V_p/V_s$  model (Model A) yields vastly superior results, both in terms of the accuracy of location and the data fit. Even the origin time is well estimated. In contrast, in the case of Model B, the 18 by 10 km error ellipse does not even encompass the true location, and the origin time is 3 seconds early. The excellent fit for Model A is somewhat surprising. The existence of significant lateral heterogeneity in crustal thickness and  $P_n$  velocity in the region (Antonenko, 1984), combined with the location of the shot outside the network, would lead one to expect less favorable results. On the one hand, it is true that the DSS profile used for the P velocity model is located between the shot point and the network, so the P structure itself may be reasonably appropriate. On the other hand, we would have expected to obtain better results with Model B, which has an S structure that is consistent with both DSS and teleseismic receiver structure results.

**Table 4**

Calculated and true locations for chemical explosions, using velocity models A and B

Explosion	Model	O.T.	Latitude	Longitude	Error ellipse (km)	True error (km)	Mean residuals
1	True	0.30	50.281	72.172			
	A	0.65	50.263	72.161	5 x 3 @ -84°	1.8	0.08 s, 1.5°
	B	-2.65	50.342	71.815	18 x 10 @ -81°	40.1	0.79 s, 2.8°
2	True	5.00	50.000	77.333			
	A	5.23	50.029	77.257	2 x 1 @ +19°	8.7	0.24 s, 2.6°
	B	5.30	50.026	77.249	7 x 3 @ +20°	9.7	0.60 s, 2.9°
3	True	0.30	50.281	72.172			
	A	1.34	50.258	72.241	18 x 9 @ -73°	7.8	0.34 s, 9.9°
	B	-2.05	50.270	71.834	22 x 13 @ -90°	37.6	0.39 s, 11.0°

For comparison, we also show the location results for explosion 3 in Table 4. Despite the overlapping teleseism, the results are nearly identical to that for explosion 1. As before, only the estimated origin time is off significantly, over 1 second late for Model A and 2 seconds early for Model B. This 3 second difference in calculated origin time for the two velocity models also mirrors the result for explosion 1. In terms of the input data, the only differences are the absence of a measured  $P_n$  azimuth for station KKL, and the fact that the data from station KSU were obtained from a high-pass-filtered version of the seismogram. Perhaps the 1 second shift in fit to the origin time between explosions 1 and 3 is due to the masking of smaller amplitude initial arrivals by the interfering teleseism.

Explosion 2 was located within the network, at a distance range from stations BAY and KKL (about 150 km) such that the first seismic phases are crustal arrivals. Unfortunately, this removes the availability of  $P_n - P_g$  arrival differences as constraints for the location, and also probably makes the identification of the first S arrival somewhat less reliable. The separation between shot 2 and station KSU is comparable to that between shot 1 and stations BAY and KKL, so mantle refracted waves are the first arrivals and a secondary crustal P phase is again observed. Azimuthal data are included in the calculations, although they were not required to produce acceptable location estimates.

Table 4 contains the location results for chemical explosion 2 for the two crustal models. The two estimated locations are nearly identical: they are both shifted 9 km west of the true location, which falls outside the error ellipse in each case. Comparison of the observed and calculated travel times indicates that the P velocity model is too slow in the upper layers, causing

the shift in location towards stations BAY and KKL. The only major difference in the results for the two models is the  $S_n$  residual at KSU, which is about 1 second late for Model A but over 3 seconds late for Model B. This suggests that the S velocities in the deeper layers of Model B are systematically too high. This is consistent with the Model B results for explosion 1, for which the calculated S arrival times were significantly early.

Since most of the regional event locations are determined using data from only two stations (BAY and KKL), it is informative to test the two-station location capability on the chemical explosions. The locations were recalculated (using Model A) excluding the data from station KSU, with little or no significant effect on the results. In the case of explosion 1, the epicenter and origin time are essentially unchanged, although the error ellipse expands by 10%. For explosion 2, the epicenter shifts 2 km westward and the origin time is 0.25 seconds later. However, the error ellipse does enlarge significantly, to 8.5 by 1.5 km. Thus in these two cases where the true event locations are known, we can derive reasonable location estimates using data from only two stations. The success of this test gives us considerable confidence in the reliability of the regional event locations discussed below, which were mostly obtained using data from two stations.

Results for the JVE compared to locations from other sources are shown above in Table 5. The difficulty in identifying clear direct S arrivals in the JVE seismograms limited its usefulness for our purposes as a master event. However the regional location results are still quite good, especially compared to the teleseismic results based on many observations.

A major product of our location effort is a catalog of well-located regional events recorded by the NRDC network. Our published locations, with two typographical errors corrected, are presented in Table 6. These locations have been used by other scientists in their studies of wave propagation in the Kazakhstan region (Chan et al., 1990; Sereno, 1990).

We had two major successes in identifying sites of industrial explosions from SPOT satellite images (Thurber et al., 1989). One was for the area around the town of Ekibastuz, north of NRDC station Bayanaul, and the other was for an area just north of Lake Balkash, south of station Karkaralinsk. In both cases, temporal changes in the appearance of surface mining or quarrying sites were detectable, lending further credence to their identification as the sources of the observed seismic events. At the time of our search through the SPOT catalog, only three other scenes were available for areas in which events were provisionally located (events a, h, and j of Thurber et al. (1989)). Only one tentative identification of an explosions site could be made - event j might be associated with a mine or quarry site at  $51^{\circ} 50' \text{ N}$ ,  $74^{\circ} 20' \text{ E}$ , located northwest of the town of Shiderty. It is possible that additional event sites could be identified, either now or in the future, if and when high-resolution satellite coverage (or maps) of the region become more available or more complete.

**Table 5**

Comparison of locations for the Joint Verification Experiment explosion in the USSR

Location source	Latitude	Longitude	Error	Uncertainty
Regional data	49.926	78.795	5 km	2 x 5 km
PDE	49.833	78.808	6 km	< 8 km
CSS	49.821	78.796	7 km	?
SPOT image	49.882	78.824	< 1 km	

**Table 6**

Catalog of 1987 two- and three-station event locations

ID	O.T. (d-h-m-s)		Latitude	Longitude	Error ellipse	Type	Area
a	1340936	16.4	50.190	74.157	26 x 3 @ -85°	blast	Karaganda
b	1350909	35.2	51.709	75.514	8 x 4 @ +74°	blast	Ekibastuz
c	1351035	0.3	49.304	72.712	3 x 2 @ +53°	blast	Karaganda
d	1410916	43.3	50.744	73.279	2 x 2 @ -80°	blast	Karaganda
e	1430849	22.7	49.275	75.738	5 x 2 @ -34°	blast	Karagayly
f	1450926	43.7	51.679	75.454	12 x 3 @ +84°	blast	Ekibastuz
g	1450956	40.9	51.743	75.316	10 x 2 @ +74°	blast	Ekibastuz
h	1460531	4.8	51.819	74.797	6 x 4 @ -85°	blast	Ekibastuz
i	1460833	26.5	51.760	75.571	15 x 6 @ -83°	blast	Ekibastuz
j	1621242	4.9	51.454	75.488	7 x 5 @ +16°	blast	Ekibastuz
k	1621250	34.3	51.677	75.525	17 x 7 @ +88°	blast	Ekibastuz
l	2340021	50.7	44.129	85.363	12 x 6 @ -69°	quake	Tien Shan
m	2390852	53.0	51.213	74.302	13 x 5 @ -26°	blast	Ekibastuz
n	2390938	34.8	46.900	77.389	6 x 3 @ +23°	blast	Balkash
o	2440344	38.8	43.808	85.948	6 x 5 @ +14°	quake	Tien Shan
p	2440908	52.0	46.924	77.241	14 x 5 @ +15°	blast	Balkash
q	2450802	10.2	51.639	75.481	12 x 6 @ -69°	blast	Ekibastuz



### 3. Conclusions and Recommendations

Starting with very little prior information, we have demonstrated the potential for a sparse seismic network of three 3-component stations, with an aperture of about 400 km, to locate events over a wide region with reasonable accuracy and precision, both for epicenter and depth. Some keys to the success of this effort have been:

- (1) the initial availability of an adequate crustal model, coupled with the inferred modest level of lateral variations in crustal structure within the array;
- (2) the availability of data from known sources (the 1987 chemical explosions) to aid the calibration of travel time models;
- (3) the existence of identifiable secondary P arrivals;
- (4) the availability of satellite imagery to confirm the source location of some events;
- (5) the ability to pursue a multifaceted approach to the investigation of the overall location problem (empirical, theoretical, observational).

We can make several specific recommendations for future efforts related to regional event location:

- (1) employ array studies to help identify and characterize secondary regional phases;
- (2) use short time windows at the onset of the P arrival for precise azimuth estimation;
- (3) obtain total coverage of the region in question by satellite imagery - including multiple sources of data (e.g., SPOT, LANDSAT).

Overall, we regard regional seismic event location to be a difficult but quite tractable problem.

#### 4. References

- Berger, J., J. N. Brune, P. A. Bodin, J. S. Gombert, D. M. Carrel, K. F. Priestley, D. E. Chavez, W. R. Walter, C. B. Archambeau, T. B. Cochran, I. L. Nersesov, M. B. Gokhberg, O. A. Stolrov, S. K. Daragen, N. D. Tarassov, and Y. A. Sutelov, A new U.S.-U.S.S.R. seismological program, *EOS, Trans. Am. Geophys. Un.*, **68**, 110-111, 1987.
- Berger, J., H. Eissler, F. L. Vernon, I. L. Nersesov, M. B. Gokhberg, O. A. Stolrov, and N. D. Tarassov, Studies of high-frequency seismic noise in eastern Kazakhstan, *Bull. Seismol. Soc. Am.*, **78**, 1744-1758, 1988.
- Bratt, S. R., and T. C. Bache, Locating events with a sparse network of regional arrays, *Bull. Seismol. Soc. Am.*, **78**, 1744-1758, 1988.
- Chan, W. W., A. Malik, and R. Baumstark, Near-crustal structure of the Soviet test site in eastern Kazakhstan, *Proc. 12th Annual DARPA/GL Seismic Res. Symp.*, pp. 18-24, 1990.
- Christoffersson, A., E. S. Husebye, and S. F. Ingate, Wavefield decomposition using ML-probabilities in modelling single-site 3-component records, *Geophys. J.*, **93**, 197-213, 1988.
- Evernden, J. F., C. B. Archambeau, and E. Cranswick, An evaluation of seismic decoupling and underground nuclear test monitoring using high-frequency seismic data, *Rev. Geophys.* **24**, 143-215, 1986.
- Given, H. K., N. T. Tarasov, V. Zhuravlev, F. L. Vernon, J. Berger and I. L. Nersesov, High-frequency seismic observations in eastern Kazakhstan, USSR, with emphasis on chemical explosion experiments, *J. Geophys. Res.*, **95**, 295-307, 1990.
- Jordan, T. H., and K. A. Sverdrup, Teleseismic location techniques and their application to earthquake clusters in the south-central Pacific, *Bull. Seismol. Soc. Am.*, **71**, 1105-1130, 1981.
- Jurkevics, A., Polarization analysis of three-component array data, *Bull. Seismol. Soc. Am.*, **78**, 1725-1743, 1988.
- Li, Y., and C. H. Thurber, Hypocenter constraint with regional seismic data: A theoretical analysis for the NRDC network in Kazakhstan, USSR, *J. Geophys. Res.*, in press, 1991.
- Magotra, N., N. Ahmed, and E. Chael, Seismic event detection and source location using single station (three-component) data, *Bull. Seismol. Soc. Am.*, **77**, 958-971, 1987.
- Ruud, B. O., E. S. Husebye, S.F. Ingate, and A. Christoffersson, Event location at any distance using seismic data from a single, three-component station, *Bull. Seismol. Soc. Am.*, **78**, 308-325, 1988.
- Sereno, T. J., Frequency-dependent attenuation in eastern Kazakhstan and implications for seismic detection thresholds in the Soviet Union, *Bull. Seismol. Soc. Am.*, **80**, 2089-2105, 1990.
- Thurber, C. H., Analysis methods for kinematic data from local earthquakes, *Rev. Geophys.*, **24**, 793-805, 1986.
- Thurber, C. H., H. Given, and J. Berger, Regional seismic event location with a sparse network: Application to eastern Kazakhstan, USSR, *J. Geophys. Res.*, **94**, 17,767-17780, 1989.

Prof. Thomas Ahrens  
Seismological Lab, 252-21  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Prof. Charles B. Archambeau  
CIRES  
University of Colorado  
Boulder, CO 80309

Dr. Thomas C. Bache, Jr.  
Science Applications Int'l Corp.  
10260 Campus Point Drive  
San Diego, CA 92121 (2 copies)

Prof. Muawia Barazangi  
Institute for the Study of the Continent  
Cornell University  
Ithaca, NY 14853

Dr. Douglas R Baumgardt  
ENSCO, Inc  
5400 Port Royal Road  
Springfield, VA 22151-2388

Prof. Jonathan Berger  
IGPP, A-025  
Scripps Institution of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Dr. Lawrence J. Burdick  
Woodward-Clyde Consultants  
566 El Dorado Street  
Pasadena, CA 91109-3245

Dr. Jerry Carter  
Center for Seismic Studies  
1300 North 17th St., Suite 1450  
Arlington, VA 22209-2308

Dr. Karl Coyner  
New England Research, Inc.  
76 Olcott Drive  
White River Junction, VT 05001

Prof. Vernon F. Cormier  
Department of Geology & Geophysics  
U-45, Room 207  
The University of Connecticut  
Storrs, CT 06268

Professor Anton W. Dainty  
Earth Resources Laboratory  
Massachusetts Institute of Technology  
42 Carleton Street  
Cambridge, MA 02142

Prof. Steven Day  
Department of Geological Sciences  
San Diego State University  
San Diego, CA 92182

Dr. Zoltan A. Der  
ENSCO, Inc.  
5400 Port Royal Road  
Springfield, VA 22151-2388

Prof. John Ferguson  
Center for Lithospheric Studies  
The University of Texas at Dallas  
P.O. Box 830688  
Richardson, TX 75083-0688

Dr. Mark D. Fisk  
Mission Research Corporation  
735 State Street  
P. O. Drawer 719  
Santa Barbara, CA 93102

Prof. Stanley Flatte  
Applied Sciences Building  
University of California  
Santa Cruz, CA 95064

Dr. Alexander Florence  
SRI International  
333 Ravenswood Avenue  
Menlo Park, CA 94025-3493

Prof. Henry L. Gray  
Vice Provost and Dean  
Department of Statistical Sciences  
Southern Methodist University  
Dallas, TX 75275

Dr. Indra Gupta  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314

Prof. David G. Harkrider  
Seismological Laboratory  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Prof. Donald V. Helmberger  
Seismological Laboratory  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Prof. Eugene Herrin  
Institute for the Study of Earth and Man  
Geophysical Laboratory  
Southern Methodist University  
Dallas, TX 75275

Prof. Bryan Isacks  
Cornell University  
Department of Geological Sciences  
SNEE Hall  
Ithaca, NY 14850

Dr. Rong-Song Jih  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314

Prof. Lane R. Johnson  
Seismographic Station  
University of California  
Berkeley, CA 94720

Dr. Richard LaCoss  
MIT-Lincoln Laboratory  
M-200B  
P. O. Box 73  
Lexington, MA 02173-0073 (3 copies)

Prof Fred K. Lamb  
University of Illinois at Urbana-Champaign  
Department of Physics  
1110 West Green Street  
Urbana, IL 61801

Prof. Charles A. Langston  
Geosciences Department  
403 Deike Building  
The Pennsylvania State University  
University Park, PA 16802

Prof. Thorne Lay  
Institute of Tectonics  
Earth Science Board  
University of California, Santa Cruz  
Santa Cruz, CA 95064

Prof. Arthur Lerner-Lam  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Dr. Christopher Lynnes  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314

Professor Peter E. Malin  
Department of Geology  
Old Chemistry Building  
Duke University  
Durham, NC 27706

Dr. Randolph Martin, III  
New England Research, Inc.  
76 Olcott Drive  
White River Junction, VT 05001

Prof. Thomas V. McEvilly  
Seismographic Station  
University of California  
Berkeley, CA 94720

Dr. Keith L. McLaughlin  
S-CUBED  
A Division of Maxwell Laboratory  
P.O. Box 1620  
La Jolla, CA 92038-1620

Prof. William Menke  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Stephen Miller  
SRI International  
333 Ravenswood Avenue  
Box AF 116  
Menlo Park, CA 94025-3493

Prof. Bernard Minster  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Prof. Brian J. Mitchell  
Department of Earth & Atmospheric Sciences  
St. Louis University  
St. Louis, MO 63156

Mr. Jack Murphy  
S-CUBED, A Division of Maxwell Laboratory  
11800 Sunrise Valley Drive  
Suite 1212  
Reston, VA 22091 (2 copies)

Prof. John A. Orcutt  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Prof. Keith Priestley  
University of Cambridge  
Bullard Labs, Dept. of Earth Sciences  
Madingley Rise, Madingley Rd.  
Cambridge CB3 0EZ, ENGLAND

Dr. Jay J. Pulli  
Radix Systems, Inc.  
2 Taft Court, Suite 203  
Rockville, MD 20850

Prof. Paul G. Richards  
Lamont Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Dr. Wilmer Rivers  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314

Prof. Charles G. Sammis  
Center for Earth Sciences  
University of Southern California  
University Park  
Los Angeles, CA 90089-0741

Prof. Christopher H. Scholz  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Thomas J. Sclero, Jr.  
Science Application Int'l Corp.  
10260 Campus Point Drive  
San Diego, CA 92121

Prof. David G. Simpson  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Dr. Jeffrey Stevens  
S-CUBED  
A Division of Maxwell Laboratory  
P.O. Box 1620  
La Jolla, CA 92038-1620

Prof. Brian Stump  
Institute for the Study of Earth & Man  
Geophysical Laboratory  
Southern Methodist University  
Dallas, TX 75275

Prof. Jeremiah Sullivan  
University of Illinois at Urbana-Champaign  
Department of Physics  
1110 West Green Street  
Urbana, IL 61801

Prof. Clifford Thurber  
University of Wisconsin-Madison  
Department of Geology & Geophysics  
1215 West Dayton Street  
Madison, WI 53706

Prof. M. Nafi Toksoz  
Earth Resources Lab  
Massachusetts Institute of Technology  
42 Carleton Street  
Cambridge, MA 02142

Prof. John E. Vidale  
University of California at Santa Cruz  
Seismological Laboratory  
Santa Cruz, CA 95064

Prof. Terry C. Wallace  
Department of Geosciences  
Building #77  
University of Arizona  
Tucson, AZ 85721

Dr. William Wortman  
Mission Research Corporation  
735 State Street  
P. O. Drawer 719  
Santa Barbara, CA 93102

OTHERS (UNITED STATES)

Dr. Monem Abdel-Gawad  
Rockwell International Science Center  
1049 Camino Dos Rios  
Thousand Oaks, CA 91360

Prof. Keiiti Aki  
Center for Earth Sciences  
University of Southern California  
University Park  
Los Angeles, CA 90089-0741

Prof. Shelton S. Alexander  
Geosciences Department  
403 Deike Building  
The Pennsylvania State University  
University Park, PA 16802

Dr. Kenneth Anderson  
BBNSTC  
Mail Stop 14/1B  
Cambridge, MA 02238

Dr. Ralph Archuleta  
Department of Geological Sciences  
University of California at Santa Barbara  
Santa Barbara, CA 93102

Dr. Jeff Barker  
Department of Geological Sciences  
State University of New York  
at Binghamton  
Vestal, NY 13901

Dr. Susan Beck  
Department of Geosciences  
Bldg. # 77  
University of Arizona  
Tucson, AZ 85721

Dr. T.J. Bennett  
S-CUBED  
A Division of Maxwell Laboratory  
11800 Sunrise Valley Drive, Suite 1212  
Reston, VA 22091

Mr. William J. Best  
907 Westwood Drive  
Vienna, VA 22180

Dr. N. Biswas  
Geophysical Institute  
University of Alaska  
Fairbanks, AK 99701

Dr. G.A. Bollinger  
Department of Geological Sciences  
Virginia Polytechnical Institute  
21044 Derring Hall  
Blacksburg, VA 24061

Dr. Stephen Bratt  
Center for Seismic Studies  
1300 North 17th Street  
Suite 1450  
Arlington, VA 22209

Michael Browne  
Teledyne Geotech  
3401 Shiloh Road  
Garland, TX 75041

Mr. Roy Burger  
1221 Serry Road  
Schenectady, NY 12309

Dr. Robert Burridge  
Schlumberger-Doll Research Center  
Old Quarry Road  
Ridgefield, CT 06877

Dr. W. Winston Chan  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314-1581

Dr. Theodore Cherry  
Science Horizons, Inc.  
710 Encinitas Blvd., Suite 200  
Encinitas, CA 92024 (2 copies)

Prof. Jon F. Claerbout  
Department of Geophysics  
Stanford University  
Stanford, CA 94305

Prof. Robert W. Clayton  
Seismological Laboratory  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Prof. F. A. Dahlen  
Geological and Geophysical Sciences  
Princeton University  
Princeton, NJ 08544-0636

Mr. Charles Doll  
Earth Resources Laboratory  
Massachusetts Institute of Technology  
42 Carleton St.  
Cambridge, MA 02142

Prof. Adam Dziewonski  
Hoffman Laboratory, Harvard Univ.  
Dept. of Earth Atmos. & Planetary Sciences  
20 Oxford St  
Cambridge, MA 02138

Prof. John Ebel  
Department of Geology & Geophysics  
Boston College  
Chestnut Hill, MA 02167

Eric Fielding  
SNEE Hall  
INSTOC  
Cornell University  
Ithaca, NY 14853

Dr. John Foley  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Prof. Donald Forsyth  
Department of Geological Sciences  
Brown University  
Providence, RI 02912

Dr. Cliff Frolich  
Institute of Geophysics  
8701 North Mopac  
Austin, TX 78759

Dr. Anthony Gangi  
Texas A&M University  
Department of Geophysics  
College Station, TX 77843

Dr. Freeman Gilbert  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California  
La Jolla, CA 92093

Mr. Edward Giller  
Pacific Sierra Research Corp.  
1401 Wilson Boulevard  
Arlington, VA 22209

Dr. Jeffrey W. Given  
SAIC  
10260 Campus Point Drive  
San Diego, CA 92121

Prof. Stephen Grand  
University of Texas at Austin  
Department of Geological Sciences  
Austin, TX 78713-7909

Prof. Roy Greenfield  
Geosciences Department  
403 Deike Building  
The Pennsylvania State University  
University Park, PA 16802

Dan N. Hagedorn  
Battelle  
Pacific Northwest Laboratories  
Battelle Boulevard  
Richland, WA 99352

Dr. James Hannon  
Lawrence Livermore National Laboratory  
P. O. Box 808  
Livermore, CA 94550

Prof. Robert B. Herrmann  
Dept. of Earth & Atmospheric Sciences  
St. Louis University  
St. Louis, MO 63156

Ms. Heidi Houston  
Seismological Laboratory  
University of California  
Santa Cruz, CA 95064

Kevin Hutchenson  
Department of Earth Sciences  
St. Louis University  
3507 Laclede  
St. Louis, MO 63103

Dr. Hans Israelsson  
Center for Seismic Studies  
1300 N. 17th Street, Suite 1450  
Arlington, VA 22209-2308

Prof. Thomas H. Jordan  
Department of Earth, Atmospheric  
and Planetary Sciences  
Massachusetts Institute of Technology  
Cambridge, MA 02139

Prof. Alan Kafka  
Department of Geology & Geophysics  
Boston College  
Chestnut Hill, MA 02167

Robert C. Kemerait  
ENSCO, Inc.  
445 Pineda Court  
Melbourne, FL 32940

William Kikendall  
Teledyne Geotech  
3401 Shiloh Road  
Garland, TX 75041

Prof. Leon Knopoff  
University of California  
Institute of Geophysics & Planetary Physics  
Los Angeles, CA 90024

Prof. L. Timothy Long  
School of Geophysical Sciences  
Georgia Institute of Technology  
Atlanta, GA 30332

Dr. Gary McCartor  
Department of Physics  
Southern Methodist University  
Dallas, TX 75275

Prof. Art McGarr  
Mail Stop 977  
Geological Survey  
345 Middlefield Rd.  
Menlo Park, CA 94025

Dr. George Mellman  
Sierra Geophysics  
11255 Kirkland Way  
Kirkland, WA 98033

Prof. John Nabelek  
College of Oceanography  
Oregon State University  
Corvallis, OR 97331

Prof. Geza Nagy  
University of California, San Diego  
Department of Ames, M.S. B-010  
La Jolla, CA 92093

Dr. Keith K. Nakanishi  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

Dr. Bao Nguyen  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Prof. Amos Nur  
Department of Geophysics  
Stanford University  
Stanford, CA 94305

Prof. Jack Oliver  
Department of Geology  
Cornell University  
Ithaca, NY 14850

Dr. Kenneth Olsen  
P. O. Box 1273  
Linwood, WA 98046-1273

Howard J. Patton  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

Prof. Robert Phinney  
Geological & Geophysical Sciences  
Princeton University  
Princeton, NJ 08544-0636

Dr. Paul Pomeroy  
Rondout Associates  
P.O. Box 224  
Stone Ridge, NY 12484

Dr. Norton Rimer  
S-CUBED  
A Division of Maxwell Laboratory  
P.O. Box 1620  
La Jolla, CA 92038-1620

Prof. Larry J. Ruff  
Department of Geological Sciences  
1006 C.C. Little Building  
University of Michigan  
Ann Arbor, MI 48109-1063



Dr. Richard Sailor  
TASC Inc.  
55 Walkers Brook Drive  
Reading, MA 01867

Dr. Susan Schwartz  
Institute of Tectonics  
1156 High St.  
Santa Cruz, CA 95064

John Sherwin  
Teledyne Geotech  
3401 Shiloh Road  
Garland, TX 75041

Dr. Matthew Sibol  
Virginia Tech  
Seismological Observatory  
4044 Derring Hall  
Blacksburg, VA 24061-0420

Dr. Albert Smith  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

Prof. Robert Smith  
Department of Geophysics  
University of Utah  
1400 East 2nd South  
Salt Lake City, UT 84112

Dr. Stewart W. Smith  
Geophysics AK-50  
University of Washington  
Seattle, WA 98195

Donald L. Springer  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

Dr. George Sutton  
Rondout Associates  
P.O. Box 224  
Stone Ridge, NY 12484

Prof. L. Sykes  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Prof. Pradeep Talwani  
Department of Geological Sciences  
University of South Carolina  
Columbia, SC 29208

Dr. David Taylor  
ENSCO, Inc.  
445 Pineda Court  
Melbourne, FL 32940

Dr. Steven R. Taylor  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

Professor Ta-Liang Teng  
Center for Earth Sciences  
University of Southern California  
University Park  
Los Angeles, CA 90089-0741

Dr. Gregory van der Vink  
IRIS, Inc.  
1616 North Fort Myer Drive  
Suite 1440  
Arlington, VA 22209

Professor Daniel Walker  
University of Hawaii  
Institute of Geophysics  
Honolulu, HI 96822

William R. Walter  
Seismological Laboratory  
University of Nevada  
Reno, NV 89557

Dr. Raymond Willeman  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Dr. Gregory Wojcik  
Weidlinger Associates  
4410 El Camino Real  
Suite 110  
Los Altos, CA 94022

Dr. Lorraine Wolf  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Prof. Francis T. Wu  
Department of Geological Sciences  
State University of New York  
at Binghamton  
Vestal, NY 13901

Dr. Gregory B. Young  
ENSCO, Inc.  
5400 Port Royal Road  
Springfield, VA 22151-2388

Dr. Eileen Vergino  
Lawrence Livermore National Laboratory  
L-205  
P. O. Box 808  
Livermore, CA 94550

J. J. Zucca  
Lawrence Livermore National Laboratory  
P. O. Box 808  
Livermore, CA 94550

GOVERNMENT

Dr. Ralph Alewine III  
DARPA/NMRO  
1400 Wilson Boulevard  
Arlington, VA 22209-2308

Mr. James C. Battis  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Dr. Robert Blandford  
AFTAC/TT  
Center for Seismic Studies  
1300 North 17th St., Suite 1450  
Arlington, VA 22209-2308

Eric Chael  
Division 9241  
Sandia Laboratory  
Albuquerque, NM 87185

Dr. John J. Cipar  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Cecil Davis  
Group P-15, Mail Stop D406  
P.O. Box 1663  
Los Alamos National Laboratory  
Los Alamos, NM 87544

Mr. Jeff Duncan  
Office of Congressman Markey  
2133 Rayburn House Bldg.  
Washington, DC 20515

Dr. Jack Evernden  
USGS - Earthquake Studies  
345 Middlefield Road  
Menlo Park, CA 94025

Art Frankel  
USGS  
922 National Center  
Reston, VA 22092

Dr. Dale Glover  
DIA/DT-1B  
Washington, DC 20301

Dr. T. Hanks  
USGS  
Nat'l Earthquake Research Center  
345 Middlefield Road  
Menlo Park, CA 94025

Paul Johnson  
ESS-4, Mail Stop J979  
Los Alamos National Laboratory  
Los Alamos, NM 87545

Janet Johnston  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Dr. Katharine Kadinsky-Cade  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Ms. Ann Kerr  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Dr. Max Koontz  
US Dept of Energy/DP 5  
Forrestal Building  
1000 Independence Avenue  
Washington, DC 20585

Dr. W.H.K. Lee  
Office of Earthquakes, Volcanoes,  
& Engineering  
345 Middlefield Road  
Menlo Park, CA 94025

Dr. William Leith  
U.S. Geological Survey  
Mail Stop 928  
Reston, VA 22092

Dr. Richard Lewis  
Director, Earthquake Engineering & Geophysics  
U.S. Army Corps of Engineers  
Box 631  
Vicksburg, MS 39180

James F. Lewkowicz  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Mr. Alfred Lieberman  
ACDA/VI-OA'State Department Bldg  
Room 5726  
320 - 21st Street, NW  
Washington, DC 20451

Stephen Mangino  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Dr. Robert Masse  
Box 25046, Mail Stop 967  
Denver Federal Center  
Denver, CO 80225

Art McGarr  
U.S. Geological Survey, MS-977  
345 Middlefield Road  
Menlo Park, CA 94025

Richard Morrow  
ACDA/VI, Room 5741  
320 21st Street N.W  
Washington, DC 20451

Dr. Carl Newton  
Los Alamos National Laboratory  
P.O. Box 1663  
Mail Stop C335, Group ESS-3  
Los Alamos, NM 87545

Dr. Kenneth H. Olsen  
Los Alamos Scientific Laboratory  
P. O. Box 1663  
Mail Stop D-406  
Los Alamos, NM 87545

Mr. Chris Paine  
Office of Senator Kennedy  
SR 315  
United States Senate  
Washington, DC 20510

Colonel Jerry J. Perrizo  
AFOSR/NP, Building 410  
Bolling AFB  
Washington, DC 20332-6448

Dr. Frank F. Pilotte  
HQ AFTAC/TT  
Patrick AFB, FL 32925-6001

Katie Poley  
CIA-ACIS/TMC  
Room 4X16NHB  
Washington, DC 20505

Mr. Jack Rachlin  
U.S. Geological Survey  
Geology, Rm 3 C136  
Mail Stop 928 National Center  
Reston, VA 22092

Dr. Robert Reinke  
WL/NTESG  
Kirtland AFB, NM 87117-6008

Dr. Byron Ristvet  
HQ DNA, Nevada Operations Office  
Attn: NVCG  
P.O. Box 98539  
Las Vegas, NV 89193

Dr. George Rothe  
HQ AFTAC/TTR  
Patrick AFB, FL 32925-6001

Dr. Alan S. Ryall, Jr.  
DARPA/NMRO  
1400 Wilson Boulevard  
Arlington, VA 22209-2308

Dr. Michael Shore  
Defense Nuclear Agency/SPSS  
6801 Telegraph Road  
Alexandria, VA 22310

Mr. Charles L. Taylor  
Phillips Laboratory - OL-AA/LWH  
Hanscom AFB, MA 01731-5000

Dr. Larry Turnbull  
CIA-OSWR/NED  
Washington, DC 20505

Dr. Thomas Weaver  
Los Alamos National Laboratory  
P.O. Box 1663, Mail Stop C335  
Los Alamos, NM 87545

Phillips Laboratory - OL-AA/ SULL  
Research Library  
Hanscom AFB , MA 01731-5000 (2 copies)

Defense Intelligence Agency  
Directorate for Scientific & Technical Intelligence  
Attn: DT1B  
Washington, DC 20340-6158

Secretary of the Air Force  
(SAFRD)  
Washington, DC 20330

AFTAC/CA  
(STINFO)  
Patrick AFB, FL 32925-6001

Office of the Secretary Defense  
DDR & E  
Washington, DC 20330

TACTEC  
Battelle Memorial Institute  
505 King Avenue  
Columbus, OH 43201 (Final Report Only)

HQ DNA  
Attn: Technical Library  
Washington, DC 20305

DARPA/RMO/RETRIEVAL  
1400 Wilson Boulevard  
Arlington, VA 22209

DARPA/RMO/Security Office  
1400 Wilson Boulevard  
Arlington, VA 22209

Phillips Laboratory  
Attn: OL-AA/XO  
Hanscom AFB, MA 01731-5000

Phillips Laboratory  
Attn: OL-AA/LW  
Hanscom AFB, MA 01731-5000

DARPA/PM  
1400 Wilson Boulevard  
Arlington, VA 22209

Defense Technical Information Center  
Cameron Station  
Alexandria, VA 22314 (5 copies)

CONTRACTORS (FOREIGN)

Dr. Ramon Cabre, S.J.  
Observatorio San Calixto  
Casilla 5939  
La Paz, Bolivia

Prof. Hans-Peter Harjes  
Institute for Geophysik  
Ruhr University/Bochum  
P.O. Box 102148  
4630 Bochum 1, FRG

Prof. Eystein Husebye  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

Prof. Brian L.N. Kennett  
Research School of Earth Sciences  
Institute of Advanced Studies  
G.P.O. Box 4  
Canberra 2601, AUSTRALIA

Dr. Bernard Massinon  
Societe Radiomana  
27 rue Claude Bernard  
75005 Paris, FRANCE (2 Copies)

Dr. Pierre Mecheler  
Societe Radiomana  
27 rue Claude Bernard  
75005 Paris, FRANCE

Dr. Svein Mykkeltveit  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY (3 copies)

FOREIGN (OTHERS)

Dr. Peter Basham  
Earth Physics Branch  
Geological Survey of Canada  
1 Observatory Crescent  
Ottawa, Ontario, CANADA K1A 0Y3

Dr. Eduard Berg  
Institute of Geophysics  
University of Hawaii  
Honolulu, HI 96822

Dr. Michel Bouchon  
I.R.I.G.M.-B.P. 68  
38402 St. Martin D'Heres  
Cedex, FRANCE

Dr. Hilmar Bungum  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

Dr. Michel Campillo  
Observatoire de Grenoble  
I.R.I.G.M.-B.P. 53  
38041 Grenoble, FRANCE

Dr. Kin Yip Chun  
Geophysics Division  
Physics Department  
University of Toronto  
Ontario, CANADA M5S 1A7

Dr. Alan Douglas  
Ministry of Defense  
Blacknest, Brimpton  
Reading RG7-4RS, UNITED KINGDOM

Dr. Roger Hansen  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

Dr. Manfred Henger  
Federal Institute for Geosciences & Nat'l Res.  
Postfach 510153  
D-3000 Hanover 51, FRG

Ms. Eva Johannisson  
Senior Research Officer  
National Defense Research Inst.  
P.O. Box 27322  
S-102 54 Stockholm, SWEDEN

Dr. Fekadu Kebede  
Geophysical Observatory, Science Faculty  
Addis Ababa University  
P. O. Box 1176  
Addis Ababa, ETHIOPIA

Dr. Tormod Kvaerna  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

Dr. Peter Marshall  
Procurement Executive  
Ministry of Defense  
Blacknest, Brimpton  
Reading FG7-4RS, UNITED KINGDOM

Prof. Ari Ben-Menahem  
Department of Applied Mathematics  
Weizman Institute of Science  
Rehovot, ISRAEL 951729

Dr. Robert North  
Geophysics Division  
Geological Survey of Canada  
1 Observatory Crescent  
Ottawa, Ontario, CANADA K1A 0Y3

Dr. Frode Ringdal  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

Dr. Jorg Schlittenhardt  
Federal Institute for Geosciences & Nat'l Res.  
Postfach 510153  
D-3000 Hannover 51, FEDERAL REPUBLIC OF GERMANY

Universita Degli Studi Di Trieste  
Facolta Di Ingegneria  
Istituto Di Miniere E. Geofisica Applicata, Trieste, ITALY

Dr. John Woodhouse  
Oxford University  
Dept of Earth Sciences  
Parks Road  
Oxford OX13PR, ENGLAND